STATISTICAL PROBES OF REIONIZATION WITH 21 CENTIMETER TOMOGRAPHY

STEVEN R. FURLANETTO,¹ MATIAS ZALDARRIAGA,^{2,3} AND LARS HERNQUIST² Received 2004 April 5; accepted 2004 June 4

ABSTRACT

We consider the degree to which 21 cm tomography of the high-redshift universe can distinguish different ionization histories. Using a new analytic model for the size distribution of H π regions that associates these ionized bubbles with large-scale galaxy overdensities, we compute the angular power spectrum and other statistical properties of the 21 cm brightness temperature during reionization. We show that the H π regions imprint features on the power spectrum that allow us to separate histories with discrete bubbles from those with partial uniform ionization (by, for example, X-rays). We also show that "double reionization" scenarios will modify the morphology of the bubbles in ways that depend on the mechanism through which the first generation of sources shuts off. If, for example, the transition occurs globally at a fixed redshift, the first generation imprints a persistent feature on the 21 cm power spectrum. Finally, we compare our model to one in which voids are ionized first. While the power spectra of these two models are qualitatively similar, we show that the underlying distributions of neutral hydrogen differ dramatically and suggest that other statistical tests can distinguish them. The next generation of low-frequency radio telescopes will have the sensitivity to distinguish all of these models and strongly constrain the history and morphology of reionization.

Subject headings: cosmology: theory — diffuse radiation — intergalactic medium — radio lines: general *Online material:* color figures

1. INTRODUCTION

Recently, a great deal of effort, both observational and theoretical, has been focused on understanding the reionization of hydrogen in the intergalactic medium (IGM) at $z \ge 6$. Several independent observational techniques offer complementary views of this landmark event. The most straightforward method is to seek extended troughs of complete Ly α absorption in the spectra of high-redshift quasars. Evidence for this Gunn & Peterson (1965) effect has been found near $z \sim 6.5$ (Becker et al. 2001; White et al. 2003). Unfortunately, the optical depth of a fully neutral medium is so high that current measurements only require a mean neutral fraction $\bar{x}_{\rm H} \gtrsim 10^{-3}$, and even this limit has been disputed (Songaila 2004). Studies of the rapidly growing ionizing background (Fan et al. 2002) and the Strömgren spheres surrounding the quasars (Wyithe & Loeb 2004) suggest larger neutral fractions but depend on detailed modeling. A second constraint arises because free electrons scatter cosmic microwave background (CMB) photons, washing out intrinsic anisotropies and generating a polarization signal (Zaldarriaga 1997). This provides an integral constraint on the reionization history; recent measurements of the CMB imply that $\bar{x}_{\rm H}$ must have been small at $z \gtrsim 14$ (Kogut et al. 2003; Spergel et al. 2003). Finally, the relatively high temperature of the Ly α forest at $z \sim 2-4$ suggests an order of unity change in $\bar{x}_{\rm H}$ at $z \leq 10$ (Theuns et al. 2002; Hui & Haiman 2003), although this argument depends on the characteristics of He II reionization (e.g., Sokasian et al. 2002).

Taken together, these three sets of observations suggest a complex ionization history extending over a large redshift in-

² Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138.

³ Jefferson Laboratory of Physics, Harvard University, Cambridge, MA 02138.

terval ($\Delta z \sim 10$). This is inconsistent with a generic picture of fast reionization (e.g., Barkana & Loeb 2001 and references therein). The results seem to indicate strong evolution in the sources responsible for reionization, and a precise measurement of the ionization history would strongly constrain early structure formation (Wyithe & Loeb 2003; Cen 2003; Haiman & Holder 2003; Sokasian et al. 2004). Unfortunately, extracting such detailed information requires new observational techniques. One of the most exciting possibilities is 21 cm tomography of the high-redshift IGM (Scott & Rees 1990; Kumar et al. 1995; Madau et al. 1997), in which one maps the distribution of neutral hydrogen on large scales through its hyperfine transition. By probing a specific spectral line, 21 cm surveys measure the time history of reionization, and unlike the Ly α forest they do not suffer from saturation problems. In fact, the signal is sufficiently weak that foregrounds from the Galaxy and a wide variety of extragalactic objects pose substantial challenges to these experiments (Oh & Mack 2003; Di Matteo et al. 2004). Fortunately, the known foregrounds all have smooth continuum spectra, which should allow frequency cleaning to high accuracy because the 21 cm signal has spectral structure on small scales (Zaldarriaga et al. 2004, hereafter ZFH04; Morales & Hewitt 2003; Cooray & Furlanetto 2004).

However, predicting the 21 cm signals from reionization has proven difficult. One method is to use numerical simulations of reionization (Ciardi & Madau 2003; Furlanetto et al. 2004a; Gnedin & Shaver 2004), but computational costs have so far limited the simulations to subtend (at best) a few resolution elements of the 21 cm maps. Analytic models of reionization can extend to larger scales but require some way to describe the complexities of structure formation and radiative transfer. The simplest approach, in which we assign each galaxy its own H II region, does a poor job of matching the large ionized bubbles found in simulations (Ciardi et al. 2003; Sokasian et al. 2003a). We have recently developed a model that associates H II regions with large-scale fluctuations in the

¹ Division of Physics, Mathematics, and Astronomy, California Institute of Technology, MC 130-33, Pasadena, CA 91125; sfurlane@tapir.caltech.edu.

density field and reproduces the major features seen in simulations (Furlanetto et al. 2004b, hereafter Paper I). This allows us, for the first time, to make quantitative predictions about the 21 cm signal at reionization. With high signal-tonoise ratio 21 cm maps, we can directly measure the distribution of H II regions in order to test our model. However, the signals are sufficiently weak that such maps will be difficult to make; fortunately, statistical measurements of the data also contain a great deal of information about reionization (ZFH04). In this paper, we use the model of Paper I to predict the 21 cm angular power spectrum for several reionization scenarios, with an emphasis on how the measurements help distinguish the crucial features of these different pictures. In \S 2 we briefly review our model for reionization. We then show how 21 cm measurements can distinguish different reionization histories in \S 3 and how they can distinguish different models of reionization in § 4. We conclude in § 5.

In our numerical calculations, we assume a Λ CDM cosmology with $\Omega_m = 0.3$, $\Omega_{\Lambda} = 0.7$, $\Omega_b = 0.046$, H = 100 h km s⁻¹ Mpc⁻¹ (with h = 0.7), n = 1, and $\sigma_8 = 0.9$, consistent with the most recent measurements (Spergel et al. 2003).

2. A MODEL FOR REIONIZATION

Recent numerical simulations (e.g., Sokasian et al. 2003a) show that reionization proceeds "inside-out" from highdensity clusters of sources to voids, at least when the sources resemble star-forming galaxies (e.g., Springel & Hernquist 2003). We therefore associate H II regions with large-scale overdensities (Barkana & Loeb 2004). We assume that a galaxy of mass m_{gal} can ionize a mass ζm_{gal} , where ζ is a constant that depends on the efficiency of ionizing photon production, the escape fraction, the star formation efficiency, and the number of recombinations. Values of $\zeta \sim 10-40$ are reasonable for normal star formation, but very massive stars can increase the efficiency by an order of magnitude (Bromm et al. 2001b). The criterion for a region to be ionized by the galaxies contained within it is then $f_{coll} > \zeta^{-1}$, where f_{coll} is the fraction of mass bound to halos above some m_{\min} . We will normally assume that this minimum mass corresponds to a virial temperature of 10⁴ K, at which point hydrogen line cooling becomes efficient. In the extended Press-Schechter model (Lacey & Cole 1993), this places a condition on the mean overdensity within a region of mass m,

$$\delta_m \ge \delta_x(m,z) \equiv \delta_c(z) - \sqrt{2}K(\zeta) \left[\sigma_{\min}^2 - \sigma^2(m)\right]^{1/2}, \quad (1)$$

where $K(\zeta) = \operatorname{erf}^{-1}(1 - \zeta^{-1})$, $\sigma^2(m)$ is the variance of density fluctuations on the scale m, $\sigma^2_{\min} = \sigma^2(m_{\min})$, and $\delta_c(z)$ is the critical density for collapse.

Paper I showed how to construct the mass function of H II regions from δ_x in an way analogous to the halo mass function (Press & Schechter 1974; Bond et al. 1991). The barrier in equation (1) is well approximated by a linear function in σ^2 , $\delta_x \approx B(m, z) = B_0 + B_1 \sigma^2(m)$. In that case, the mass function has an analytic expression (Sheth 1998):

$$m\frac{dn}{dm} = \sqrt{\frac{2}{\pi}\frac{\bar{\rho}}{m}} \left| \frac{d\ln\sigma}{d\ln m} \right| \frac{B_0}{\sigma(m)} \exp\left[-\frac{B^2(m,z)}{2\sigma^2(m)} \right], \quad (2)$$

where $\bar{\rho}$ is the mean density of the universe. Equation (2) gives the comoving number density of H II regions with masses in the range (m, m + dm). The crucial difference be-

tween this formula and the standard Press-Schechter mass function arises from the fact that δ_x is a (decreasing) function of *m*. The barrier is more difficult to cross as one goes to smaller scales, which gives the bubbles a characteristic size. In contrast, the barrier used in constructing the halo mass function is independent of mass, yielding the usual power-law behavior at small masses. The characteristic scale depends primarily on $\bar{x}_{\rm H}$.

Paper I also showed how to construct the power spectrum of the quantity $\psi = x_{\rm H}(1 + \delta)$. We explicitly computed the correlation function of the H II regions ξ_{xx} in terms of the mass function in equation (2), and we computed the density correlation function $\xi_{\delta\delta}$ and cross correlation between density and ionization fraction $\xi_{x\delta}$ using the halo model (e.g., Cooray & Sheth 2002). We combined these correlation functions to get ξ_{ψ} , the correlation function of ψ ,

$$\xi_{\psi} = \xi_{xx}(1 + \xi_{\delta\delta}) + \bar{x}_{H}^{2}\xi_{\delta\delta} + \xi_{x\delta}(2\bar{x}_{H} + \xi_{x\delta}).$$
(3)

ZFH04 showed how to convert this to the three-dimensional power spectrum P_{ψ} and then to the (observable) angular power spectrum of the 21 cm brightness temperature δT_b , essentially by doing a Fourier transform. Here

$$\delta T_b \approx 23\psi \left(\frac{T_S - T_{\rm CMB}}{T_S}\right) \left(\frac{\Omega_b h^2}{0.02}\right) \\ \times \left[\left(\frac{0.15}{\Omega_m h^2}\right) \left(\frac{1+z}{10}\right) \right]^{1/2} \,\mathrm{mK}, \tag{4}$$

with T_S the spin temperature of the gas. In all of the calculations here, we assume $T_S \gg T_{\text{CMB}}$, which should be reasonable even relatively early in reionization (see § 2 of ZFH04). We further assume perfect frequency resolution in computing the angular power spectrum; our results therefore do not depend on the experimental setup. The effects of finite bandwidth are described in ZFH04 and Paper I. We refer the reader to these papers for more details on our approach. Note that these methods to construct the angular power spectrum of δT_b can be applied to any model of the H II regions.

3. COMPLEX REIONIZATION

Paper I examined the power spectra of standard reionization scenarios in which all the ionizations are confined to discrete H II regions. We showed that the power spectrum evolves rapidly throughout reionization, allowing one to reconstruct the ionization history in such a simple scenario. The thin curves in Figure 1 show the evolution of the angular power spectrum in a "standard" reionization scenario with $\zeta = 40$. As reionization progresses, the H II regions (obviously) fill up more space, adding power as in equation (3), and grow rapidly, shifting that power to larger scales. Figure 1 shows that the resulting power spectrum develops a prominent feature at the characteristic bubble scale late in reionization. The strength and angular size of the feature allow us to measure how the bubble topology evolves through reionization. Interestingly, the bubble size distribution and its power spectrum P_{xx} (on linear scales) are only weakly dependent on ζ for a fixed $\bar{x}_{\rm H}$ (Paper I). To first order, increasing or decreasing ζ merely translates the distribution in time. This in turn implies that the location and strength of the bubble feature change only insofar as the angular diameter distance depends on



Fig. 1.—The 21 cm fluctuation amplitude as a function of redshift. The thick curves have $\zeta = 12$ plus a uniformly ionized component, while the thin curves have $\zeta = 40$. The ionization fraction $\bar{x}_{\rm H}(z)$ is identical in the two models. The curves have $\bar{x}_{\rm H} = 0.96$ (*dotted curve*), $\bar{x}_{\rm H} = 0.8$ (*dashed curve*), and $\bar{x}_{\rm H} = 0.26$ (*solid curve*). [See the electronic edition of the Journal for a color version of this figure.]

redshift.⁴ Thus, the thin curves in Figure 1 give the approximate power spectrum for *any* single reionization episode. We refer the reader to Figure 7 of Paper I for an explicit example of how much the power spectrum changes with ζ . Here we extend our model in order to consider the effects of more complicated reionization processes.

3.1. A Partially Ionized IGM

We first consider how well 21 cm observations can distinguish a model with all ionizations inside of H II regions from one with a uniform component, such as may happen through ionization by X-rays (Oh 2001; Venkatesan et al. 2001; Ricotti & Ostriker 2004; Madau et al. 2004) or decaying particles (Sciama 1982; Hansen & Haiman 2004). Suppose that the IGM has a uniform ionization fraction $\bar{x}_u(z)$: on top of this uniform level are some variations in the neutral fraction owing to isolated H II regions. In this case, the condition $f_{coll} > \zeta^{-1}(1 - \bar{x}_u)$ replaces the barrier in our model: each galaxy can produce a larger ionized bubble with the same number of ionizing photons. However, rather than varying from zero to unity, we have $0 < x_H < (1 - \bar{x}_u)$. This damps the fluctuations from the bubbles, requiring the replacements $\xi_{xx} \rightarrow (1 - \bar{x}_u)^2 \xi_{xx}$ and $\xi_{x\delta} \rightarrow$ $(1 - \bar{x}_u)\xi_{x\delta}$. Otherwise, our formalism is unchanged.

Figure 1 contrasts such a case to the "standard" history described above. The thick curves have $\zeta = 12$ and set $\bar{x}_u(z)$ to force the total neutral fraction to match in the two cases. The uniform component is thus responsible for 70% of the ionizations. When $\bar{x}_H \approx 1$, the two curves are nearly identical

because the bubble feature is intrinsically weak. However, while the bubbles rapidly imprint a feature on the power spectrum if $\bar{x}_u = 0$, uniform ionization strongly damps the power from the H II regions. As a result, in such a scenario the 21 cm fluctuations trace those of the density (normalized by $1 - \bar{x}_u$), at least until reionization is almost complete. Consequently, the amplitude of δT_b is suppressed by 25%–50% on scales near the characteristic bubble size, even when \bar{x}_H is large. We find that, unless reionization is near completion, the bubbles need to provide a majority of the ionizations in order for them to have an appreciable effect on the power spectrum. This provides a clear test for whether Strömgren spheres from ultraviolet photons or smoother ionization from some other source dominate the morphology of reionization, even if the survey is unable to make high signal-to-noise ratio maps.

3.2. Double Reionization

Recently, a number of theoretical models for reionization have attempted to reconcile the CMB and quasar data by postulating an early generation of sources with high ionizing efficiency (most often because they contain massive, metal-free stars) along with a self-regulation mechanism that switches to normal star formation with a lower ionizing efficiency (Wyithe & Loeb 2003; Cen 2003; Haiman & Holder 2003; Sokasian et al. 2004). Such scenarios can lead to a plateau in the ionized fraction or even "double" reionization, in which ionized phases bracket a mostly neutral period. In this section, we explore the the signatures that such scenarios imprint on the 21 cm power spectrum through some simple toy models.

We consider an early generation of sources with $(\zeta_1, m_{\min,1})$ and a later one with $(\zeta_2, m_{\min,2})$. In the initial phase only the first generation is important, so the ionization simply follows the model of Paper I. We then end this generation according to one of two conditions. The first choice shuts down these sources once the universe reaches some uniform level of ionization at redshift z_1 . For example, the early phase could be caused by X-rays or decaying particles, as described in § 3.1. A more realistic scenario could be the following. Suppose that the first generation consisted of massive, metal-free stars; the natural self-regulation condition halts the formation of these stars when the metallicity in collapsed objects passes some threshold (Bromm et al. 2001a; Mackey et al. 2003; Yoshida et al. 2004). These sources will stop forming soon after they first appear in any given region (or, in other words, soon after the region is ionized for the first time) but they will continue to form in the remaining neutral regions. Thus, the first generation will persist until the entire universe has been ionized. Our toy model makes the simplification that the first generation was able to keep the universe ionized until the transition occurs; we implicitly neglect recombinations within regions that were ionized before z_1 .

While the second generation is still rare, recombination in the IGM will be the dominant effect. Such an era is obviously easy to identify with 21 cm tomography because it is accompanied by a corresponding increase in $\bar{x}_{\rm H}$ and hence in the signal (Furlanetto et al. 2004a). If the recombination rate were uniform throughout the IGM, we would have the power spectrum $P_{\psi} = \bar{x}_{\rm H}^2 P_{\delta\delta}$. Because the recombination rate is a function of the density, we also get fluctuations in the neutral fraction that amplify δT_b . In the limit in which the elapsed time is smaller than the recombination time, we would have $P_{xx} \approx [1 - \bar{x}_i(z)/\bar{x}_i(z_1)]^2 P_{\delta\delta}$, where \bar{x}_i is the mean ionized fraction. Unfortunately, $x_{\rm H}(\delta, z)$ is a nonlinear function of density and time if we include recombinations, so the detailed evolution must

⁴ This is not strictly true if nonlinear corrections to the density field cannot be ignored on the relevant scales (which can happen if reionization occurs at $z \leq 7$). However, even in this case both the shape and amplitude of the 21 cm power spectrum are demonstrably different from the underlying matter power spectrum.

await numerical simulations. Given the weak large-scale fluctuations present at high redshifts, we do not expect recombinations to amplify the power by more than a few tens of percent, and it should not imprint prominent features into the power spectrum. (Note that allowing extra recombinations in regions that were ionized before z_1 would increase this amplification factor.) In our formalism, we can implement this scenario using the results of § 3.1, with the extra wrinkle that \bar{x}_u can decrease with cosmic time.

Another prescription shuts off the first generation before those sources can fully ionize the universe. At z_1 , the universe has developed a patchwork of H II regions but they have not yet overlapped completely. This choice is appropriate if the selfregulation mechanism is not local. For example, the buildup of an ultraviolet background in the Lyman-Werner bands could halt H₂ cooling and shut off star formation in small halos (Haiman et al. 1997), or an X-ray background could heat the universe and raise the Jeans mass. In this model, the first generation imprints a set of ionized bubbles, within which most of the second-generation sources grow (because both appear in the same overdense regions). The bubbles grow only slowly until the total number of ionizations from the second generation become comparable to that of the first; after this point the evolution approaches the normal behavior. If we neglect recombinations within H II regions, the barrier can be set up as follows. Consider a region of mass *m* at $z_2 < z_1$; as in Paper I, we need the condition for sources inside this region to ionize it. The number of ionizations is simply the sum of those from both generations. Thus, we find that the excursion set barrier $\delta_x(m)$ will be the solution of

$$1 = (\zeta_1 - \zeta_2) \operatorname{erfc} \left\{ \frac{\delta_c(z_1) - \delta_x(m)}{\sqrt{2[\sigma^2(m_{\min,1}) - \sigma^2(m)]}} \right\} + \zeta_2 \operatorname{erfc} \left\{ \frac{\delta_c(z_2) - \delta_x(m)}{\sqrt{2[\sigma^2(m_{\min,2}) - \sigma^2(m)]}} \right\}.$$
(5)

Here the complementary error functions are the fraction of collapsed gas above the mass thresholds at the two redshifts. This is a more complicated expression than in Paper I, but δ_x is still approximately linear in σ^2 . With the new barrier, the formalism from Paper I carries over without further modification. We could crudely incorporate recombinations into this model by decreasing ζ_1 as redshift decreases. This is not exactly correct, because it would cause the bubbles to shrink rather than recombining at all radii. However, this may not be a serious problem because, according to our prescription, both types of sources will primarily form within the same highly biased regions. Even a relatively small number of secondgeneration ionizing sources could be enough to halt recombinations within the bubble unless the recombination time is much smaller than the Hubble time. In any case, the main feature of this model is that the first-generation bubbles imprint a particular scale on the power spectrum. If recombination is significant (but not complete) and the secondgeneration sources reside in partially ionized bubbles, their own H II regions will grow faster than normal and will quickly reach this scale, at which point their expansion will slow. If, on the other hand, recombination is nearly complete in these regions, all evidence of the first generation disappears and the single-generation description of Paper I would apply.

Figure 2 contrasts the power spectra of three scenarios for double reionization. In the solid curves, the first sources uni-



FIG. 2.—The 21 cm power spectrum in three different models of "double reionization." The solid curves assume that the universe has $\bar{x}_u = 0.5$, the long-dashed curves assume $z_1 = 18$ with $\zeta_1 = 500$, and the short-dashed curves assume $z_1 = 18$ with $\zeta_1 = 105$ and m_{\min} 10 times smaller than our default value. In all cases, $\zeta_2 = 12$. The thin curves are for z = 16 ($\bar{x}_H = 0.46$) and the thick curves are for z = 12 ($\bar{x}_H = 0.27$). [See the electronic edition of the Journal for a color version of this figure.]

formly ionize the IGM to $\bar{x}_u = 0.5$ (or, if we neglect amplification due to recombinations, the IGM has recombined to this point after a phase of full ionization). In the other models, we end the first generation at $z_1 = 18$ when $\bar{x}_{\rm H} = 0.5$; the dotted and dashed curves correspond to two different sets of parameters for the first generation of sources. In all cases, the second generation has $\zeta_2 = 12$ and ionizes $\sim 4\%$ (z = 16) and $\sim 23\%$ (z = 12) of the IGM. Note that we have ignored recombinations after z_1 in order to isolate the effects of different prescriptions. The most obvious feature of the second type of transition is that the bubble size remains approximately constant with redshift. This contrasts sharply with scenarios in which a single set of sources is responsible for reionization, for which the bubble scale changes rapidly over $\Delta z \sim 1$ (Paper I). We suggest that this plateau in the bubble scale is a clear indication that the transition between generations occurs rapidly and in a nonlocal manner. It is, however, difficult to distinguish between the two scenarios containing relic bubbles. As described in Paper I, this is because the scale of the H II regions depends only on the behavior of the large-scale density fluctuations and is fixed almost entirely by $\bar{x}_{\rm H}$. In the uniform model, the amplitude of δT_b is significantly smaller and the bubble features are suppressed, just as in § 3.1. Note also that in this case the bubble feature grows rapidly, because the H II regions expand into a preionized medium. This stands in stark contrast to the dashed curves. While including recombination could increase the overall amplitude of the uniform model by a small amount, it would not affect the bubble feature.

4. OUTSIDE-IN REIONIZATION

In Paper I and in § 3, we have examined how the parameters of the ionizing sources change the 21 cm power spectrum; however, we have always worked within the model described

in \S 2. It is easy to imagine many other models for reionization, and we now consider whether the 21 cm signal can discriminate between them. For example, if recombinations control reionization, we expect low-density regions to be ionized first (Miralda-Escudé et al. 2000). This should certainly be true on small scales: within the H II regions, low-density regions will remain ionized, while denser regions will stay neutral until the ionizing background increases. Miralda-Escudé et al. (2000) implement this by assuming that all gas elements below a fixed density threshold are ionized first, with the threshold increasing as reionization progresses. This model is meant to apply on relatively small scales (i.e., smaller than the mean separation between sources). However, the general idea could also apply to larger scales if, for example, rare but bright sources are responsible for reionization. In such a scenario, voids could be ionized first because they are not able to recombine during the periods without nearby, active sources. We refer to this as "outside-in" reionization, because the dense regions (where galaxies sit) are ionized last. A simple way to describe such a process is to assume that regions below a fixed mean overdensity $\delta_v < 0$ are ionized. We can then construct the mass function of the ionized bubbles through the excursion set formalism; the result is identical to the Press & Schechter (1974) halo mass function except that $|\delta_v|$ replaces the collapse threshold.⁵ We fix δ_v by requiring the fraction of mass in ionized regions with $m > \zeta m_{\min}$ (i.e., larger than an H II region around the smallest allowed galaxy) to equal ζf_{coll} . This prescription is undoubtedly an oversimplification, but it nonetheless serves to illustrate the importance and testability of our basic assumptions. The only additional ingredient we need to construct the power spectrum is the bias of ionized regions. With the above prescription, we have

$$b_{v}(m,z) = -\left(1 + \frac{\delta_{v}^{2}/\sigma^{2} - 1}{|\delta_{v}|}\right).$$
 (6)

This is the same expression as for the halo bias in the Press-Schechter model (Mo & White 1996), except that the bias is negative because the number density of voids increases in *underdense* regions.⁶

We compare the outside-in angular power spectrum to the inside-out model of Paper I in Figure 3. We have normalized the two models to have identical $\bar{x}_{\rm H}(z)$. We see that, until relatively late in reionization, the fluctuation amplitude and the characteristic bubble scale are similar in the two models. This is because both models associate H II regions with largescale fluctuations in the density field. For the same reason, the void size distribution is only weakly dependent on ζ for a fixed $\bar{x}_{\rm H}$ and the difference between the two models is essentially independent of when reionization occurs. The only difference is that the void model barrier δ_v is independent of scale but δ_x increases nearly linearly with σ^2 in the inside-out model. As a result, the void model has somewhat more ionized gas in smaller regions and a somewhat smaller characteristic scale; however, these differences may not be robust given the approximate methods used to construct the size distribution.

On the other hand, the similarity shows explicitly that, regardless of the detailed model, the characteristic bubble size is large (and in the ideal range for 21 cm tomography) so long as the sizes of H II regions are determined by fluctuations in the large-scale density field. A more fundamental difference between the two models is that the void model has significantly more large-scale power, at least until shot noise from the bubbles begins to dominate at $z \sim 13$.⁷ By associating ionized regions with underdensities, the void model *amplifies* largescale power. In contrast, inside-out reionization washes out

large-scale power by ionizing overdensities first.

curves). In both cases, $\zeta = 40$. The curves are $\bar{x}_{\rm H} = 0.96$ (z = 18, dotted

curve), $\bar{x}_{\rm H} = 0.8$ (z = 15, *dashed curve*), and $\bar{x}_{\rm H} = 0.26$ (z = 12, *solid curve*). [See the electronic edition of the Journal for a color version of this figure.]

Moreover, as emphasized in Paper I, the power spectrum does not uniquely determine the statistics of the δT_b field. Although the density fluctuations are nearly Gaussian on the relevant scales, the H II regions are not. We now consider the pixel distribution function (PDF) in the two models; i.e., the fraction of spherical "pixels" of a given mass m_{pix} that have brightness temperature δT_b . We showed how to compute this for the insideout model in Paper I.⁸ The same technique applies to the void model once we modify the excursion set barrier appropriately.⁹ We show some example PDFs in Figure 4; they assume $m_{pix} = 10^{13} M_{\odot}$ or an angular resolution $\sim 2.'7$ at z = 13. The figure explicitly demonstrates that although both reionization models increase the variance in δT_b , they do so in qualitatively different ways. For inside-out reionization, highdensity regions are highly ionized and δT_b has a maximum



⁵ In principle, we should include the "void-in-cloud" process described by Sheth & van de Weygaert (2004) in order to remove those voids that have been consumed by collapsed objects. However, the regime in which we are interested has $|\delta_v| \ll \delta_c$, so this process turns out to have no effect on our results.

⁶ With this prescription, we can have the cross correlation between a halo and a void $\xi_{hv} < -1$, which is unphysical. In these situations we set $\xi_{hv} = -1$; fortunately, this has a negligible effect on our results.

 $^{^{7}}$ As described in Paper I, our model probably begins to break down at this point because the internal structure of the H II regions becomes important.

⁸ Note that we do not include overlap with large bubbles in the results shown here, so these PDFs cannot be directly compared with observations; see Paper I.

⁹ In this case $x_{\rm H}(\delta)$ is a monotonic function, so the resulting PDF is nonsingular.



FIG. 4.—PDF (*top*) and cumulative distribution function $P(>\delta T_b)$ (*bottom*) in our three models, assuming $\zeta = 40$ and $m_{\text{pix}} = 10^{13} M_{\odot}$. The solid, dashed, and dotted curves are for the inside-out model of Paper I, the outside-in void model, and a uniformly ionized IGM, respectively. In the bottom panel, the three sets of curves are for z = 20 ($\bar{x}_{\text{H}} = 0.98$), z = 16 ($\bar{x}_{\text{H}} = 0.88$), and z = 13 ($\bar{x}_{\text{H}} = 0.5$), from right to left. We only show the PDFs for the latter two cases. [See the electronic edition of the Journal for a color version of this figure.]

value δT_{max} on any given scale. The pixels tend to cluster strongly around this maximum value because δT_h is a weak function of density in the relevant range; the PDF has large variance because the mean $\overline{\delta T} < \delta T_{\text{max}}$, thanks to fully ionized regions. For outside-in reionization, on the other hand, highdensity pixels remain mostly neutral, while the peak and the low-density tail get stretched to small δT_b . The result is not so far from a Gaussian, but $x_{\rm H}(\delta)$ has dramatically increased the variance. The bottom panel of Figure 4 shows the cumulative distribution functions $P(>\delta T_b)$ for three redshifts, which is often useful in statistical tests. Note that at early times, when $\bar{x}_{\rm H}$ is small, the distributions closely resemble each other, but significant features appear even when the ionized fraction is only ~10%. Also, the inside-out model has $p(\delta T_b = 0) \sim 0.2$ at z = 13, because the characteristic bubble size is near m_{pix} . Figure 4 shows the necessity of identifying useful tests for non-Gaussianity in the signal. We intend to explore such tests in the future.

5. DISCUSSION

We have examined what 21 cm tomography of the highredshift universe can teach us about the ionization history. Using the reionization model described in Paper I, we predicted the angular power spectra of the brightness temperature δT_b in several qualitatively different reionization scenarios. We have shown that 21 cm measurements offer a powerful probe of the

reionization process, even without high signal-to-noise ratio maps. Most importantly, the angular power spectrum evolves throughout reionization, with a bubble feature appearing and shifting to larger scales as more and more gas is ionized. The location and strength of this feature is determined by the largescale density field and depends only weakly on when reionization occurs. Moreover, nonstandard ionization histories have measurable effects on the power spectrum. For example, we showed that the "bubble" feature in the power spectrum disappears if a substantial fraction of the ionizations are due to hard photons that can travel large distances from their source. Also, we contrasted the predictions of two different prescriptions for the transition between multiple generations of sources. In one, the transition occurs in an approximately uniformly ionized medium. In the other, the transition occurs while the original H II regions are still growing. We showed that these different prescriptions lead to qualitatively different power spectra, even long after the first sources have shut off, because the first generation bubbles can imprint a persistent feature on the power spectrum. This could be especially important if the first phase of reionization ends at redshifts inaccessible to the next generation of radio telescopes, because of either radio frequency interference or simple noise considerations (note that the background is approximately proportional to $[1+z]^{2.7}$ for these measurements). Finally, we contrasted our "inside-out" reionization model with predictions for an "outside-in" model in which voids are ionized before dense regions. While the power spectrum of these models differ in detail, we showed that the underlying distributions of pixel brightness temperature for a given angular resolution are dramatically different. We thus suggest that more sophisticated statistical measures of the data will be able to strongly constrain the underlying model of reionization.

Of course, we have made a number of simplifying assumptions in all of these models. Most notably, we have ignored recombinations, radiative transfer, and the anisotropic distribution of sources; we discuss these issues in more detail in Paper I. Addressing these complications in a satisfactory manner will probably require numerical simulations. However, none are likely to alter our conclusions: 21 cm measurements contain an unprecedented wealth of information about the timing and morphology of reionization. Our simple model shows that the qualitative statistical features of the signal can be related to the source and IGM properties. The next generation of low-frequency radio telescopes, such as the Primeval Structure Telescope,¹⁰ the Low Frequency Array,¹¹ and the Square Kilometer Array,¹² should be able to measure these statistical properties (ZFH04) and place strong constraints on the reionization era.

This work was supported in part by NSF grants ACI AST 99-00877, AST 00-71019, AST 0098606, and PHY 0116590, and NASA ATP grants NAG5-12140 and NAG5-13292 and by the David and Lucille Packard Foundation Fellowship for Science and Engineering.

¹¹ See http://www.lofar.org for details on LOFAR.

¹² See http://www.skatelescope.org for details on the SKA.

REFERENCES

- Barkana, R., & Loeb, A. 2001, Phys. Rep., 349, 125
- _____. 2004, ApJ, 609, 474
- Becker, R. H., et al. 2001, AJ, 122, 2850
- Bond, J. R., Cole, S., Efstathiou, G., & Kaiser, N. 1991, ApJ, 379, 440
- Bromm, V., Ferrara, A., Coppi, P. S., & Larson, R. B. 2001a, MNRAS, 328, 969
- Bromm, V., Kudritzki, R. P., & Loeb, A. 2001b, ApJ, 552, 464
- Cen, R. 2003, ApJ, 591, L5
- Ciardi, B., & Madau, P. 2003, ApJ, 596, 1
- Ciardi, B., Stoehr, F., & White, S. D. M. 2003, MNRAS, 343, 1101
- Cooray, A., & Furlanetto, S. R. 2004, ApJ, 606, L5
- Cooray, A., & Sheth, R. 2002, Phys. Rep., 372, 1

¹⁰ See http://astrophysics.phys.cmu.edu/~jbp/past6.pdf for details on PAST.

- Di Matteo, T., Ciardi, B., & Miniati, F. 2004, MNRAS, submitted (astro-ph/ 0402322)
- Fan, X., et al. 2002, AJ, 123, 1247
- Furlanetto, S. R., Sokasian, A., & Hernquist, L. 2004a, MNRAS, 347, 187
- Furlanetto, S. R., Zaldarriaga, M., & Hernquist, L. 2004b, ApJ, 613, 1 (Paper I)
- Gnedin, N. Y., & Shaver, P. A. 2004, ApJ, 608, 611
- Gunn, J. E., & Peterson, B. A. 1965, ApJ, 142, 1633
- Haiman, Z., & Holder, G. P. 2003, ApJ, 595, 1
- Haiman, Z., Rees, M. J., & Loeb, A. 1997, ApJ, 476, 458 Hansen, S. H., & Haiman, Z. 2004, ApJ, 600, 26
- Hui, L., & Haiman, Z. 2003, ApJ, 596, 9
- Kogut, A., et al. 2003, ApJS, 148, 161
- Kumar, A., Padmanabhan, T., & Subramanian, K. 1995, MNRAS, 272, 544
- Lacey, C., & Cole, S. 1993, MNRAS, 262, 627
- Mackey, J., Bromm, V., & Hernquist, L. 2003, ApJ, 586, 1
- Madau, P., Meiksin, A., & Rees, M. J. 1997, ApJ, 475, 429
- Madau, P., et al. 2004, ApJ, 604, 484
- Miralda-Escudé, J., Haehnelt, M., & Rees, M. J. 2000, ApJ, 530, 1
- Mo, H. J., & White, S. D. M. 1996, MNRAS, 282, 347
- Morales, M. F., & Hewitt, J. 2003, ApJ, submitted (astro-ph/0312437)
- Oh, S. P. 2001, ApJ, 553, 499
- Oh, S. P., & Mack, K. J. 2003, MNRAS, 346, 871

- Press, W. H., & Schechter, P. 1974, ApJ, 187, 425
- Ricotti, M., & Ostriker, J. P. 2004, MNRAS, 352, 547
- Sciama, D. W. 1982, MNRAS, 198, 1P
- Scott, D., & Rees, M. J. 1990, MNRAS, 247, 510
- Sheth, R. K. 1998, MNRAS, 300, 1057
- Sheth, R. K., & van de Weygaert, R. 2004, MNRAS, 350, 517
- Sokasian, A., Abel, T., & Hernquist, L. 2002, MNRAS, 332, 601
- Sokasian, A., Abel, T., Hernquist, L., & Springel, V. 2003a, MNRAS, 344, 607 Sokasian, A., Yoshida, N., Abel, T., Hernquist, L., & Springel, V. 2004,
- MNRAS, 350, 47 Songaila, A. 2004, AJ, 127, 2598
- Spergel, D. N., et al. 2003, ApJS, 148, 175
- Springel, V., & Hernquist, L. 2003, MNRAS, 339, 312
- Theuns, T., et al. 2002, ApJ, 567, L103
- Venkatesan, A., Giroux, M. L., & Shull, J. M. 2001, ApJ, 563, 1
- White, R. L., Becker, R. H., Fan, X., & Strauss, M. A. 2003, AJ, 126, 1
- Wyithe, J. S. B., & Loeb, A. 2003, ApJ, 588, L69
- 2004, Nature, 427, 815
- Yoshida, N., Bromm, V., & Hernquist, L. 2004, ApJ, 605, 579
- Zaldarriaga, M. 1997, Phys. Rev. D, 55, 1822
- Zaldarriaga, M., Furlanetto, S. R., & Hernquist, L. 2004, ApJ, 608, 622 (ZFH04)